

in E

already monitored plant responses to recent climate change that include (1) colonization, as altitudinal and latitudinal displacements occur (Parmesan and Yohe, 2003) and (2) adaptation, such as spring-time advances in phenological processes (Hughes, 2000). Currently, mortality through increased attacks by pests and diseases following drought stress is putting some ecosystems at risk (Woods et al., 2005; Hicke et al., 2006). Because of the anticipated rates of climate change, population re-locations through dispersal and colonization are expected to

first is through environmentally-induced epigenetic change and the second is through stochastic epimutations (Verhoeven and Preite, 2014).

Environmentally-induced epigenetic changes are direct responses to environmental stresses that are heritable across generations and distinct from common environmental effects such as on gametes or seeds subject to the same environment as the parent. Such effects have been shown for salt-stressed invasive knotweeds (Richards et al., 2008), environmentally-stressed dandelions (Verhoeven et al., 2010) and pathogen-infected *Arabidopsis* (Downen et al., 2012). Because these epigenetic

responses to select for stress-tolerant genotypes are unlikely to be rapid enough to protect populations against extinction in the face of extreme stresses. However, the potential for epigenetic responses to stress may provide the phenotypic variation necessary to sustain populations during events that could push plants past threshold tolerance levels. We now

- drought–fire interactions. *Proc. Natl. Acad. Sci. U.S.A.* 111, 6347–6352. doi: 10.1073/pnas.1305499111
- Bräutigam, K., Vining, K. J., Lafon-Placette, C., Fossdal, C. G., Mirouze, M., Marcos, J. G., et al. (2013). Epigenetic regulation of adaptive responses of forest tree species to the environment. *Ecol. Evol.* 3, 399–415. doi: 10.1002/ece3.461
- Castonguay, E., and Angers, B. (2012). The key role of epigenetics in the persistence of asexual lineages. *Genet. Res. Int.*

- Rapp, R. A., and Wendel, J. F. (2005). Epigenetics and plant evolution. *New Phytol.* 168, 81–91. doi: 10.1111/j.1469-8137.2005.01491.x
- Richards, C. L., Bossdorf, O., Muth, N. Z., Gurevitch, J., and Pigliucci, M. (2006). Jack of all trades, master of some? On the role of phenotypic plasticity in plant invasions. *Ecol. Lett.* 9, 981–993. doi: 10.1111/j.1461-0248.2006.00950.x
- Richards, C. L., Schrey, A. W., and Pigliucci, M. (2012). Invasion of diverse habitats by few Japanese knotweed genotypes is correlated with epigenetic differentiation. *Ecol. Lett.* 15, 1016–1025. doi: 10.1111/j.1461-0248.2012.01824.x
- Richards, C. L., Walls, R. L., Bailey, J. P., Parameswaran, R., George, T., and Pigliucci, M. (2008). Plasticity in salt tolerance traits allows for invasion of novel habitat by Japanese knotweed s. l. (*Fallopia japonica* and *F. bohemica*, Polygonaceae). *Am. J. Bot.* 95, 931–942. doi: 10.3732/ajb.2007364
- Richards, E. J. (2006). Inherited epigenetic variation—revisiting soft inheritance. *Nat. Rev. Genet.* 7, 395–401. doi: 10.1038/nrg1834
- Savidan, Y. (2010). Apomixis: genetics and breeding.