

positive phonotaxis, turning towards the sound source. Conversely,

literature indicates that bats (a natural predator of crickets) emit echolocation signals at frequencies around 18 kHz (Moiseff et al., 1978). Night time induces flying behavior in crickets, and when hearing predatorial bat ultrasounds, crickets perform negative phonotaxis and fly away from the sound (Moiseff et al., 1978). Past experiments indicate that crickets have a dominant ear. When bat ultrasounds are presented between 20 degrees to the left or right either directly in front of the cricket or behind it, the animal is not able to localize the direction of the stimulus. Instead, the cricket will consistently turn to either the left or right direction, meaning that crickets have a dominant ear (Nolen and Hoy 1986).

The auditory organs of the crickets are located on the foreleg, and auditory afferents protrude into a central nervous system hub located inferior to the prothoracic ganglion. In healthy crickets, auditory neurons from both sides of the body send projections to the respective side within the prothoracic ganglion as they extend their projections into the brain. When a cricket is deafferented (meaning it's foreleg is cut off, and thus its auditory organs as well), a compensatory neuroplasticity phenomenon occurs. Neuroplasticity refers to the reorganization of neurons, in this case after injury. Acutely post deafferentation, the cricket loses the bilateral communication from the ear to the brain, losing the ability to localize sound sources. Through an unknown mechanism, the auditory neurons from the deafferented side are able to cross the previously respective side in the prothoracic ganglion and make synapses with the contralateral side (Fischer et al., 2018). While auditory inputs are only received from the intact ear, axonal projections proceed up both sides of the prothoracic ganglion into the brain for processing and potentially localization.

Experiments this summer aimed to figure out where compensatory plasticity occurs and how it is beneficial to the survival of crickets. The experiment was studied through a behavioral lens, characterizing the flight behavior of crickets in novel environments, evaluating whether the cricket regains localization abilities after the auditory system neuroplasticity occurs. I started analysis on past behavior experiments done in the Horch Lab using the `BehavioralCide` and `epLabCut`, and this quantitative

analysis is still ongoing.

Preliminary results from the past experiments have suggested that crickets do not lose their localization abilities, as turn angles in response to stimuli have

decreased in response to deafferentation.

