

**Investigating the impact of an RNA-binding protein on hyphal tip localization of an mRNA  
in *Candida albicans*  
Alajjah Rubianes, Class of 2024**

*Candida albicans* is an opportunistic pathogenic fungus that can enter the bloodstream & internal organs, which can lead to life-threatening infections for immunocompromised people [1, 2]. Infection rate is linked to the transformation of budding yeast cells to hyphal cells [3]. Hyphal

**Faculty Mentor: McBride**

**Funded by the Kibbe Science Fellowship**

**References:**

1. Candida infections of the mouth, throat, and esophagus | Fungal Diseases | CDC. 2021 Feb 25. <https://www.cdc.gov/fungal/diseases/candidiasis/thrush/index.html>
2. Buchacz K, Lau B, Jing Y, Bosch R, Abraham AG, Gill MJ, Silverberg MJ, Goedert JJ, Sterling TR, Althoff KN, et al. Incidence of AIDS-Defining Opportunistic Infections in a Multicohort Analysis of HIV-infected Persons in the United States and Canada, 2000–2010. *The Journal of Infectious Diseases*. 2016;214(6):862–872. doi:[10.1093/infdis/jiw085](https://doi.org/10.1093/infdis/jiw085)
3. Lo HJ, Köhler/F3 1bm6z K K